

Post-operative Instructions for Periodontal Surgery

I. Right after your Surgery

- 1. Following surgery, proper rest is important in aiding healing and reducing the possibility of discomfort, bleeding, or other complications.
- 2. Do not rinse for 12 hours. After 12 hours you may gently rinse with warm water after meals.
- 3. Do not eat for 2 hours after surgery.
- 4. During the next 3 days, avoid excessive exertion of any type (golf, tennis, jogging, mowing lawn, heavy lifting, yoga, etc.). You may follow your regular daily activities provided they do not require a great deal of physical exertion.
- 5. You might consider using an extra pillow to elevate your head above chest level.
- 6. During healing, you may experience tooth sensitivity with temperature changes (particularly to cold).

 This is usually temporary and should gradually diminish and eliminate itself in approximately 6-8 weeks.
- 7. While your mouth is numb, be careful to not bite your cheek, lip, or tongue. The numbness should subside within 2-4 hours.

II. Medications

- 1. Take medications as prescribed. Follow instructions for all medication accurately.
- 2. For postoperative comfort have your prescriptions filled as soon as possible.

III. Diet

- 1. It is extremely important that you maintain your regular diet as much as possible in order to promote proper healing. Avoid chewing with the teeth where surgery was performed.
- 2. Avoid hot liquids or foods. Allow for liquid to cool to room temperature.
- 3. Avoid eating hard foods that require a lot of chewing pressure, spicy foods, nuts, chips, or popcorn, etc.
- 4. Drink plenty of fluids the first 48 hours after surgery.
- 5. Avoid alcohol.

IV. Oral Hygiene

- 1. Brush parts of mouth on which surgery was **not** performed.
- 2. Floss your teeth but not in the area of surgery until advised otherwise.
- 3. Rinse your mouth carefully after eating.

V. You may Experience:

- 1. A slight amount of blood seepage may occur, giving your saliva a red color. Do not be alarmed. Direct pressure with wet gauze or a wet tea bag for 20 minutes should control bleeding. If the seepage persists, call Dr. Wang.
- 2. Swelling and bruising may occur, and is normal-this may last several days.
- 3. If you have pain and the medicine prescribed for you does not give you relief, please contact Dr. Wang. Office: (603) 224-9474.